

1 Original Article

2 **The macroecology of animal versus wind pollination: ecological factors are more important**
3 **than historical climate stability**

4 André Rodrigo Rech¹, Bo Dalsgaard², Brody Sandel³, Jesper Sonne², Jens-Christian Svenning³,
5 Naomi Holmes⁴, Jeff Ollerton^{4*}

6

7 *¹Universidade Federal dos Vales do Jequitinhonha e Mucuri (UFVJM), Diamantina, Minas Gerais,*
8 *Brazil; ²Center for Macroecology, Evolution and Climate, Natural History Museum of Denmark,*
9 *University of Copenhagen, Copenhagen, Denmark; ³Section for Ecoinformatics and Biodiversity,*
10 *Department of Bioscience, Aarhus University, Aarhus, Denmark; ⁴Landscape and Biodiversity*
11 *Research Group, School of Science and Technology, University of Northampton, Northampton, UK.*

12 Short running-title: The macroecology of pollination

13 * Corresponding author. Email: Jeff.Ollerton@northampton.ac.uk

14

1 **Abstract**

2 **Background:** The relative frequency of wind and animal pollinated plants is non-randomly
3 distributed across the globe and numerous hypotheses have been raised for the greater occurrence of
4 wind pollination in some habitats and towards higher latitudes. To date, however, there has been no
5 comprehensive global investigation of these hypotheses.

6 **Aims:** Investigating a range of hypotheses for the role of biotic and abiotic factors as determinants
7 of the global variation in animal versus wind pollination.

8 **Methods:** We analysed 67 plant communities ranging from 70° North to 34° South. For these we
9 determined habitat type, species richness, insularity, topographic heterogeneity, current climate and
10 Late-Quaternary climate change. The predictive effects of these factors on the proportion of wind-
11 and animal-pollinated plants were tested using correlations, ordinary least squares (OLS) and
12 logistic regression analyses with information-theoretic model selection.

13 **Results:** The proportion of animal-pollinated plant species was positively associated with plant
14 species richness and current temperature. Furthermore, in forest, animal pollination was positively
15 related to precipitation. Historical climate was only weakly and idiosyncratically correlated with
16 animal pollination.

17 **Conclusion:** Results were consistent with the hypothesised reduced chance for wind-transported
18 pollen reaching conspecific flowers in species-rich communities, fewer constraints on nectar
19 production in warm and wet habitats, and reduced relative effectiveness of wind dispersal in humid
20 areas. There was little evidence of a legacy of historical climate change affecting these patterns.

21 **Keywords:** abiotic, biotic, community ecology, forest, mutualism, open vegetation, pollen
22 dispersal, precipitation, species richness, temperature.

23

24

1 **Introduction**

2 Pollination is a fundamental ecological process that is crucial for the functioning of most terrestrial
3 ecosystems (Kearns et al. 1998; Ollerton et al., 2011; Lever et al., 2014). Plants achieve pollen transfer
4 among conspecifics using biotic (animal) and abiotic (wind and, rarely, water) pollen vectors. Wind
5 pollination occurs in about 20% of angiosperm families and most groups of gymnosperms (Ackerman
6 2000), an estimated 12.5% of all angiosperm species are wind pollinated (Ollerton et al. 2011). Wind
7 pollination is a derived condition that has evolved independently more than 60 times within otherwise
8 animal-pollinated clades (Ackerman 2000; Friedman and Barrett 2009; Hu et al. 2012), with
9 occasional reversions to biotic pollination (e.g. Wragg and Johnson 2011).

10 Wind pollination is currently understood to be a reproductive strategy that evolves when
11 pollinators become unpredictable or unavailable (Barrett 1996; Culley et al. 2002; Friedman and
12 Barrett 2009). Supporting this idea, the proportion of wind-pollinated plants is higher in the
13 temperate zone (Culley et al. 2002; Ollerton et al. 2011), where climate is less stable seasonally
14 (Cramer et al. 1999) and over longer glacial–interglacial time scales (Sandel et al. 2011). In
15 temperate regions usually < 80% of the plant species within a community are animal pollinated,
16 whereas animal pollination is more common in the tropics where on average ≥ 90 % of all co-
17 occurring plants are pollinated by animals (Whitehead 1969; Regal 1982; Ollerton et al. 2011).
18 Although this latitudinal pattern is already known, no study has quantitatively assessed the possible
19 factors related to the global variation of community-level differences in wind *versus* animal
20 pollination (Barrett 1996; Schemske et al. 2009; Ollerton et al. 2011).

21 The efficiency of wind and animal pollination may depend on the biotic and abiotic
22 environment where plants occur. A number of testable hypotheses for geographic variation in wind
23 *versus* animal pollination have been put forward (Whitehead 1969; Culley et al. 2002). For instance,
24 extensive, dense and species-rich forests with high canopy and low wind speeds are supposed to be

1 unfavourable places for wind pollination because the chance of wind transporting pollen to a
2 conspecific should be rather low (Whitehead 1969; Regal 1982; Ollerton et al. 2006, 2011; Culley
3 et al. 2002). Climate may also affect the efficiency of wind pollination. High humidity should make
4 the cytoplasm of the pollen grains to collapse or become heavier and cause them to clump, reducing
5 pollen dispersal distance (Whitehead 1969; Niklas 1985). Additionally, high temperature and
6 precipitation may increase productivity and support more nectar production for animal pollination.

7 Thus, both mean annual temperature and precipitation should correlate positively with the
8 proportion of animal pollination. On the other hand, high temperature and precipitation seasonality
9 may promote wind pollination as pollinator abundances will fluctuate through the year, becoming
10 less reliable pollen vectors (Regal 1982), unless pollination is in synchrony with these seasonal
11 fluctuations in pollinator abundances, thereby causing no measurable effect on pollination mode
12 (McKinney et al. 2012). Finally, in topographically flat environments such as coastal plains wind
13 pollination could be favoured due to more constant and fast winds. Likewise wind pollination could
14 be more frequent on islands because wind-pollinated plants are independent of pollinator
15 colonisation to maintain reproduction (Kühn et al. 2006; Crawford et al. 2011).

16 Paleoclimate has been linked to contemporary ecological patterns, such as species
17 distributions and endemism (Svenning and Skov 2007; Cárdenas et al. 2011; Sandel et al. 2011;
18 Kissling et al. 2012), and the structure of interactions within communities of plants and pollinators
19 (Dalsgaard et al., 2011, 2013; Groom et al., 2014). Past climatic instability may have broken up
20 mutualistic associations between plant species and their pollinating animals (Dalsgaard et al., 2011,
21 2013, McKinney *et al.*, 2012) that may have favoured wind pollination and could be related to the
22 contemporary composition of plant communities.

23 Here, we use a global dataset to describe (1) latitudinal patterns in pollination mode, and (2)
24 the relationships between putative contemporary and historical factors, and the proportion of animal

1 *versus* wind pollinated species in plant communities around the world. Specifically, following on
2 from the discussion of abiotic and biotic factors that other authors have presented, we test the
3 following hypotheses:

4 1. Wind pollination is less frequent in closed forest habitats compared to more open habitats,
5 such as grasslands.

6 2. Animal pollination is more frequent in areas with higher mean annual temperature and
7 precipitation.

8 3. Conversely, wind pollination is more frequent in regions with lower temperature and
9 greater seasonality of precipitation.

10 4. Animal pollination is more frequent in communities with high plant species richness.

11 5. Wind pollination is more frequent in topographically flat environments.

12 6. Animal pollination is less frequent on islands.

13 7. Wind pollination is more frequent in those areas of the world that have experienced
14 greater climatic instability during the Quaternary.

15

16 **Materials and methods**

17 *Animal versus wind pollination datasets*

18 The dataset was taken from 82 published and unpublished community-based assessments of the
19 proportion of wind and animal pollinated species. These studies are censuses of all native
20 angiosperm species in flower in a given community at a specific time. Gymnosperms were not
21 considered by most of the published studies, hence we excluded gymnosperms from all
22 communities. The latitudes ranged from 34° S to 70° N (Figure 1). When the same geographic
23 coordinates were attributed to communities within the same vegetation type by the original authors,
24 we pooled them together and used the mean of pollination mode proportion as a descriptor of the

1 proportion of animal pollination for those coordinates. Doing this, we reduced the dataset to 67 data
2 points for the spatial analysis (Appendix Table S1).

3 In some of the published community studies no species list was available, just data on
4 the proportions of wind and animal pollinated species observed, preventing us from attributing
5 genera or families to the data analysed by the original authors. For those studies having a species
6 list, pollination mode at the species level was highly correlated with higher taxa, i.e. genera and
7 families (Table 1). This does not mean that species within the same genus have always the same
8 pollination mode. It implies that proportions are constant across taxonomic levels and therefore any
9 level may be used in the analysis. For the published studies we followed the information provided
10 by the original authors as to the pollination mode (wind or animal) of a given species. For our own
11 field observations, flowers were assessed in terms of presence or absence of: mass pollen release
12 when an inflorescence was shaken, reduced corolla, and feathery stigmas (wind pollination); or
13 large and colourful flowers with scent or nectar, and the presence and behaviour of potential flower
14 visitors (animal pollination) (Figure 2; see also Table 1 in Friedman and Barrett 2009). We paid
15 particular attention to species with small, dull flowers, e.g. some tropical trees, where only the
16 presence of scent, nectar, and/or insect visitation distinguished biotic from abiotic pollination.

17 The proportion of animal-pollinated plant species could be underestimated because some plant
18 species that possess wind pollination traits can also be visited and sometimes pollinated by animals
19 (Figueredo and Sazima 2000). There might also be cases of obligatory self-pollinating and non-
20 sexually reproducing species coded as wind or animal pollinated, though this should be a relatively
21 minor and not spatially structured bias and, hence, should not affect our results (Ollerton et al. 2011).
22 Similarly to Ollerton et al. (2011), we took a community-level approach as we were interested in the
23 ecological question of how large a proportion of plant species in terrestrial communities are animal
24 or wind pollinated, and the possible factors related to these proportions. However, first, we measured

1 the phylogenetic signal for pollination mode by means of Phylogenetic Eigenvector Regression
2 (PVR) considering the species lists where plants were identified up to the species level (Diniz-Filho
3 et al. 2012). Species lists were available for 56 plant communities, including 1689 genera.
4 Phylogenetic information was extracted at genus level from Phylomatic (R20120829 stored tree,
5 Webb and Donoghue 2005) and the tree generated by using Phylocom (Webb et al. 2008). PVR was
6 calculated by using a principal coordinate analysis (PCoA) of the phylogenetic distance matrix and
7 selecting the 50 first axes. Phylogenetic signal was obtained by the logistic regression coefficient (R^2)
8 between the selected phylogenetic eigenvectors and the binomial pollination mode (Diniz-Filho et al.
9 2012). Phylogenetic signal significance was tested by measuring PVR on 1000 randomisations of the
10 pollination mode in the phylogeny. The analysis was carried out by using the package PVR for R
11 (Santos et al. 2013). Since phylogenetic signal of pollination mode was very low (PVR for 50
12 eigenvectors = 0.0411; $P = 0.47$; $DF = 50$; Deviance = 50.2) no taxonomic correction was required.
13 The low phylogenetic signal probably came out of many independent origins of wind pollination and,
14 due to many genera having one pollination mode also showing a minority of species possessing the
15 other pollination mode.

16

17 *Correlates of pollination mode*

18 For each site, we extracted variables of contemporary and historical climate conditions
19 hypothesised to affect the degree of wind *versus* animal pollination. Current climate descriptors
20 included mean annual temperature (MAT), mean annual precipitation (MAP), temperature
21 seasonality (MAT seasonality) and precipitation seasonality (MAP seasonality) extracted at 2.5 arc-
22 minute resolution (approximately 16 km²) from the WorldClim dataset (www.worldclim.org;
23 Hijmans et al. 2005). As different climate change metrics may capture different effects of climate
24 change (Garcia et al. 2014), we considered two measures of historical climate stability: (1) the

1 velocity of mean annual temperature (MAT velocity) and that of mean annual precipitation (MAP
2 velocity) between the Last Glacial Maximum (LGM) and the present; and (2) the anomaly of mean
3 annual temperature (MAT anomaly) and mean annual precipitation (MAP anomaly) between the
4 LGM and the present (current minus LGM precipitation/temperature – i.e. positive values indicate
5 wetter or warmer contemporary conditions, respectively). Velocities reflect the speed of climate
6 change irrespective of the direction, integrating macroclimatic changes and local spatial topoclimate
7 gradients (Loarie et al. 2009; Sandel et al. 2011), whereas anomalies reflect the direction of climate
8 change. Paleoclimate estimates were obtained from the CCSM3 model (Collins et al. 2006; Otto-
9 Bliesner et al. 2006), and statistically downscaled to 2.5 arc-minute resolution to match the current
10 climate data (Hijmans et al. 2005).

11 In addition, we determined potentially important non-climate descriptors of local
12 conditions at each site, including whether the site was on the mainland (coded as 0) or on an island
13 (1), topographic heterogeneity, regional proportion of tree cover, plant species richness, and
14 whether the vegetation was forest (0; n = 16) or open vegetation type (1; grassland, scrubland, and
15 other low and open vegetation types, n = 51). Topographic heterogeneity was calculated as the
16 range of elevation values observed in a $0.2^\circ \times 0.2^\circ$ window (ca. 20 km \times 20 km) centered on the
17 sampling location, using the WorldClim 1-km DEM (Farr et al. 2007). The regional proportion of
18 tree cover was calculated as the average tree cover observed within a window of 64 km x 64 km
19 around each site, based on the Vegetation Continuous Fields product applied to MODIS data
20 (Hansen et al. 2003).

21

22 *Statistical analyses*

23 For all analyses, plant species richness, MAT anomaly, precipitation and temperature
24 velocities were \log_{10} -transformed; contemporary mean annual precipitation and topography were

1 square-root transformed; the proportion of regional tree cover was arcsine square-root transformed.
2 The proportion of animal pollinated plant species in each community was arcsine square-root
3 transformed for ordinary least squares (OLS) regression, but was left untransformed for logistic
4 regression. All other variables were left untransformed.

5 First, the latitudinal pattern in pollination mode was tested with a quadratic regression
6 between latitude and the proportion of animal pollination. Second, for all vegetation types, and
7 separately for open vegetation types and forest, we tested the univariate associations between all
8 predictor variables and between the proportion of animal pollinated plant species and all predictor
9 variables (Appendix Table S2). This identified predictor variables that were strongly inter-
10 correlated. For all further analyses, we excluded temperature seasonality as it was strongly
11 correlated with mean annual temperature ($r = -0.73$, $P < 0.001$, $n = 67$), and we excluded the
12 proportion of regional tree cover as it was unavailable for three island datasets and strongly
13 correlated with contemporary precipitation ($r = 0.73$, $P < 0.001$, $n = 64$; see Appendix Table S2).
14 Due to the strong positive relationship between precipitation and the proportion of regional tree
15 cover, we adjusted our above hypothesis of a unidirectional positive effect of precipitation on
16 animal pollination. For forest, we still expected a positive correlation between precipitation and the
17 proportion of animal pollination. For open vegetation types, on the other hand, we expected a less
18 strong (or even zero) effect of precipitation due to the possible opposite effects of increasing
19 precipitation (favouring animal pollination) and associated increasing fragmentation of open
20 vegetation communities surrounded by increasing amounts of forests as precipitation and regional
21 tree cover coincide (increased fragmentation favouring wind pollination).

22 To test the study hypotheses, we used ordinary least squares (OLS) and logistic
23 regression, conducting information-theoretic model selection, as outlined by Diniz-Filho et al.
24 (2008). In doing so, we tested the effect of both measures of historical climate stability, i.e.

1 temperature/precipitation velocity and anomaly; the anomaly models generally performed the best
2 (highest R^2 and lowest AIC), so we focus the discussion on these models (but we also report models
3 including velocity). First, we fitted models with all combinations of the explanatory variables:
4 forest versus open vegetation, insularity, plant species richness, contemporary precipitation,
5 contemporary temperature, precipitation seasonality, topography and historical climate stability as
6 measured by temperature and precipitation anomaly. Furthermore, due to the possible different
7 effect of precipitation in open and forest vegetation, we included an interaction term between
8 precipitation and forest versus open vegetation. We then identified minimum adequate models
9 (MAMs) among these as any model with $\Delta AIC_c < 2$ ($n = 67$, 1023 alternative models; Table 2).
10 When using temperature and precipitation velocity as a measure of climate stability, we excluded
11 topography as it is strongly correlated with velocities (Appendix S2), i.e. when using velocity as our
12 historical climate variable we compared 511 models. We were unable to estimate velocities for one
13 island data point, therefore in models using velocities there is one data point fewer.

14 Standardised regression coefficients are reported for both a multi-model average
15 regression model based on weighted w_i and the MAMs (Diniz-Filho et al. 2008). We tested whether
16 significant ($P < 0.05$) positive spatial autocorrelation remained in OLS model residuals, tested using
17 10 distance classes and applying a permutation test with 10,000 iterations. As no spatial
18 autocorrelation was found in our anomaly models that performed the best, we did not use spatial
19 models. All correlations and OLS regression analyses were conducted using Spatial Analysis in
20 Macroecology 4.0 (Rangel et al. 2010). Logistic model selection analyses were carried out in R (R
21 Development Core Team, 2014) using the ‘MuMIn’ package (Barton 2015).

22

23 **Results**

1 The proportion of animal-pollinated plants in a community decreased with latitude (Figure 1). This
2 pattern appeared to be largely symmetrical across the Equator, though there was a geographical bias
3 to the distribution of the studies, and more data from Africa and Asia, in particular, are needed to
4 confirm the observed pattern.

5 The models to assess the relative importance of the factors tested as determinants of
6 the proportion of animal pollination showed that overall the explanatory power was considerably
7 higher for current climate variables (ca. $32\% < R^2 < 36\%$) than for species richness (ca. $4\% < R^2 <$
8 5%) and historical climate variables (ca. $0\% < R^2 < 2\%$; Table 2 and Appendix S3). The proportion
9 of animal pollination in both simple univariate correlations and in regression models was strongly
10 and positively associated with current temperature and species richness (Figure 3; Table 2;
11 Appendix S3). Furthermore, the interaction term between precipitation and vegetation type was
12 included in the best-fit regression model (Table 2; Appendix S3), indicating that precipitation
13 correlated strongly with the proportion of animal pollination in forest (Figure 3; Appendix S2).
14 There were only weak associations with topography and temperature anomaly and velocity (Table
15 2, Appendix S2). In addition to explaining little variation in pollination mode, topography and
16 historical temperature anomaly/velocity associated positively with animal pollination in regression
17 models (Table 2), but were unrelated in simple univariate correlations (Table S1). In logistic, but
18 not in OLS, regression islands were found to have a lower proportion of animal pollination.
19 Seasonality, precipitation anomaly, and precipitation velocity were not statistically significant in
20 simple univariate correlations nor were they included in any MAMs (Table 2 and Appendix S3).

21

22 **Discussion**

23 The proportion of plant species using animal versus wind pollination in a given community is non-
24 random with respect to latitude, animal pollination being especially dominant in the tropics (Figure

1 1), confirming previous studies, but with a much larger and geographically widespread data set. We
2 found that differences in the proportion of animal pollinated species were associated mainly with
3 current climate (Figure 3). Notably, temperature was overall positively related to the prevalence of
4 animal pollination (partly supporting hypothesis 2), and in forests animal pollination was more
5 frequent in areas with higher precipitation (supporting both hypothesis 1 and, in part, hypotheses 2
6 and 3). However, the ‘more wind pollination with greater precipitation seasonality’ aspect of
7 hypothesis 3 was not supported, perhaps due to the different nature of seasonality in the lowland
8 tropics (wet, dry) compared with the temperate or high altitude zones (cold and warm). In addition
9 to current climate, we found that communities with greater local plant species richness have a
10 higher proportion of animal pollination (supporting hypothesis 4). On the other hand, topography
11 had little effect on frequency of pollination mode, not supporting hypothesis 5. Kühn et al. (2006)
12 showed that wind pollination in Germany is more frequent in the flat landscapes of north relative to
13 other parts of the country, although it increases also toward the southern Alpine uplands. This
14 suggests that the effect of topography is scale- and context-dependent, which may explain its weak
15 effect in our global analysis. Insularity also had only a minor influence on pollination-mode, with
16 islands tending to have lower proportions of animal-pollination, but only when also taking climate
17 into account (Table 2), which only partly supports hypothesis 6. In contrast to the strong
18 associations with current factors (Table 2; Appendix S2), historical climate was associated only
19 weakly and idiosyncratically with animal pollination, varying independently when using regression
20 models or simple univariate correlations (Table 2 and Appendix S2), and therefore not supporting
21 hypothesis 7. Overall, this suggests that the relative proportions of different pollination modes are
22 more linked to current climate and ecological factors than to historical legacies.

23 Although these results support the greater influence of contemporary ecological
24 factors over historical ones, it is not straightforward to determine the exact mechanism responsible

1 for this pattern. For example, current precipitation was strongly related to the percentage of regional
2 tree cover, hence, animal pollination is associated with both high precipitation and forest habitat.
3 Wet forests clearly offer poor conditions for pollen dispersal, but it is difficult to disentangle
4 whether this is because of mechanical restrictions imposed by dense vegetation or because air
5 humidity decreases pollen dispersability, or both (Whitehead 1969; Niklas 1985). In open areas,
6 especially in the tropics, the distribution of plant individuals tends to be sparse and floral resources
7 less abundant. For example, an estimate of nectar energy available per hectare in the Atlantic rain
8 forest in Brazil showed that forests produced twice as much floral energy as the nearby open
9 ‘restinga’ (coastal scrub) vegetation (Fonseca 2013, Fonseca et al. 2015). Moreover, the distribution
10 of an Andean plant species was associated with a progressive disruption of its animal pollination
11 mutualism accompanied by the gradual decrease in precipitation (Chalcoff et al. 2012). This
12 illustrates the potential role of vegetation structure and precipitation in determining the functioning
13 of animal-dependent mutualisms, but disentangling the roles of correlated factors is a major
14 challenge for macroecology.

15 Interestingly, current climate may play a similar role for seed dispersal as there is also
16 a pattern of higher importance of animal dispersers in rain forests, whereas in dry forests wind and
17 self-dispersal are more common (Howe and Smallwood 1982; Jordano 2000 and references
18 therein). For instance, current precipitation patterns are largely correlated with proportion of
19 endozoochory across the Atlantic forest of Brazil, animals being more important dispersers in the
20 wetter areas (Almeida-Netto et al. 2008), although large-scale studies are not available to confirm
21 this hypothesis. One tentative joint explanation for the lower frequency of animal-pollinated and
22 animal-dispersed plants at drier sites could be the higher metabolic costs of producing nectar and
23 fleshy fruit, as was originally proposed only for fruit (Wilson et al. 1989; Almeida-Netto et al.
24 2008). Associated with humidity, high temperature may also increase nectar production and

1 therefore influence animal pollination, although, as temperature and temperature seasonality were
2 found to be negatively correlated, we cannot rule out the possibility that seasonality promotes wind
3 pollination (Regal 1982).

4 Another factor positively associated with animal pollination was plant species
5 richness. Other trophic interactions between plants and animals respond to the bottom-up effect of
6 plant species richness (Kissling et al. 2007; Scherber et al. 2010); for instance, in the Neotropics the
7 diversity of pollinating birds and bats is associated with their food plant species richness (Fleming
8 2005; Kissling et al. 2007). Meanwhile, as plant species richness increases, the density of
9 conspecifics per unit area tends to decrease (Comita et al. 2010), which may also reduce the
10 efficiency of pollen dispersion by wind (Whitehead 1969; Regal 1982). Plant species richness may
11 also play an important role in sustaining pollinator communities (Ebeling et al. 2008; Dorado and
12 Vázquez 2014) via two main mechanisms: (1) more plant species per unit area may ensure a more
13 predictable and diverse food supply through complementarity (Waser and Real 1979; Rathcke 1983;
14 Blüthgen and Klein 2011; Yang et al. 2013); and (2) more plant species also increases species
15 redundancy within functional groups ('biodiversity insurance effect'), reducing the extinction risk
16 of functionally specialised interactions (Bartomeus et al. 2013; Fründ et al. 2013). For instance,
17 manipulative experiments have shown that high functional diversity of plants enhances the number
18 of pollinators, and the functional diversity of pollinators increases individual plant fitness and
19 community persistence over time (Fontaine et al. 2006; Albrecht et al. 2012).

20

21 **Conclusions**

22 We tested a range of previously proposed hypotheses and found that a subset of contemporary
23 climate and ecological processes are the factors more strongly related to the global pattern of animal
24 than to wind pollination at the community level. The deduced limited influence of historical climate

1 instability on the contemporary prevalence of animal and wind pollination is in contrast with
2 previous findings of high importance of historical factors on the structuring of plant-pollinator
3 interaction networks (Dalsgaard et al., 2011, 2013; Martín González et al. 2015) and the distribution
4 of animal and plant life on earth (Svenning and Skov 2007; Cárdenas et al. 2011; Sandel et al. 2011;
5 Kissling et al. 2012), but is in accordance with studies of the organisation of plant-frugivore
6 interaction networks (Schleuning et al. 2014). In future studies, it would be interesting to examine if
7 other measures of historical processes, e.g. geological differences or historical measures of plant
8 migration rates, influence plant reproduction. The large scale drivers of plant reproductive ecology
9 will be better understood with similar studies examining also the role of both historical and
10 contemporary factors on other processes, such as plant reproduction via autogamy and seed
11 dispersal. In addition, to improve the certainties of the conclusions presented here will require
12 additional sampling effort in Africa, Asia, Russia and oceanic islands, where so few data have been
13 collected on plant reproductive ecology at the community level.

14

15 **Acknowledgements**

16 Special thanks to Dr. Andrew Moldenke, who shared with us his great personal community
17 assessments dataset, and to all the researchers who helped us during field work, pollination mode
18 confirmation, community assessments and/or with extra data to perform this study. ARR was
19 supported by FAPESP (2009/54491-0); CAPES (PDSE 0197/12-2); CNPq (140912/2010-0);
20 Unicamp (FAEPEX 519292) and Santander Universities. BD and JS acknowledge the Danish
21 National Research Foundation for funding the Center for Macroecology, Evolution and Climate
22 (grant-number DNRF96). JCS was supported by the European Research Council (ERC-2012-StG-
23 310886-HISTFUNC). JO was supported by FAPESP; The Royal Society; The Leverhulme Trust;
24 The British Ecological Society; Church and Co. PLC; EVOLINK; The Percy Sladen Memorial Fund;

1 The Royal Entomological Society and the University of Northampton. NH was supported by the
2 British Exploring Society.

3

4

5 **References**

6

7 Ackerman JD. 2000. Abiotic pollen and pollination, ecological, functional, and evolutionary
8 perspectives. *Plant Systematics and Evolution* 222: 167-185.

9 Albrech M, Schmid B, Hautier Y, Müller CB. 2012. Diverse pollinator communities enhance plant
10 reproductive success. *Proceedings of the Royal Society B* 279: 4845-4852. Almeida-Neto M,
11 Campassi F, Galetti M, Jordano P, Oliveira-Filho A. 2008. Vertebrate dispersal syndromes
12 along the Atlantic forest: broad-scale patterns and macroecological correlates. *Global Ecology*
13 *and Biogeography* 17: 503–513.

14 Barton K. 2015. MuMIn: Multi-Model Inference. R package version 1.13.4. Available at:
15 <http://cran.r-project.org/package=MuMIn>.

16 Barrett SCH. 1996. The reproductive biology and genetics of island plants. *Philosophical*
17 *Transactions of the Royal Society Ser. B.* 351: 725–733.

18 Bartomeus I, Park MG, Gibbs J, Danforth BN, Lakso AN, Winfree R. 2013. Biodiversity ensures
19 plant–pollinator phenological synchrony against climate change. *Ecology Letters* 11: 1331–
20 1338.

21 Blüthgen N, Klein AM. 2011. Functional complementarity and specialisation: Why biodiversity is
22 important in plant-pollinator interactions. *Basic and Applied Ecology* 12: 282–291.

- 1 Cárdenas ML, Gosling WD, Sherlock SC, Poole I, Pennington RT, Mothes P. 2011. The response of
2 vegetation on the Andean flank in western Amazonia to pleistocene climate change. *Science*
3 331: 1055-1058.
- 4 Chalcoff VR, Aizen MA, Ezcurra C. 2012. Erosion of a pollination mutualism along an
5 environmental gradient in a south Andean treelet, *Embothrium coccineum* (Proteaceae). *Oikos*
6 121: 471-480.
- 7 Collins WD, Bitz CM, Blackman ML et al. 2006. The Community Climate System Model version 3
8 (CCSM3). *Journal of Climate* 19: 2122-2143.
- 9 Comita LS, Muller-Landau HC, Aguilar S, Hubbell SP. 2010. Asymmetric density dependence
10 shapes species abundances in a tropical tree community. *Science* 329: 330-332.
- 11 Cramer W, Kicklighter DW, Bondeau A, Iii BM, Churkina G, Nemry B, Ruimy A, Schloss AL. 1999.
12 Comparing global models of terrestrial net primary productivity (NPP): overview and key
13 results. *Global Change Biology* 5: 1–15.
- 14 Crawford DJ, Anderson GJ, Bernardello G. 2011. The reproductive biology of Island plants. *The*
15 *Biology of Island Floras* (ed. by D. Bramwell and J. Caujapé-Castells), pp. 11-36, Cambridge
16 University Press.
- 17 Culley TM, Weller SG, Sakai AK. 2002. The evolution of wind pollination in angiosperms. *Trends*
18 *in Ecology and Evolution* 17: 361-369.
- 19 Dalsgaard B, Magård E, Fjeldså J. et al. 2011. Specialization in plant-hummingbird networks is
20 associated with species richness, contemporary precipitation and quaternary climate-change
21 velocity. *PLoS One* 6: e25891.
- 22 Dalsgaard B, Trøjelsgaard K, González AMM. et al. 2013. Historical climate-change influences
23 modularity and nestedness of pollination networks. *Ecography* 36: 1331-1340.

- 1 Diniz-Filho JAF, Bini LM, Rangel TF, Morales-Castilla I, Olalla-Tárraga MÁ, Rodríguez MÁ,
2 Hawkins BA. 2012. On the selection of phylogenetic eigenvectors for ecological analyses.
3 *Ecography* 35: 239.
- 4 Diniz-Filho JAF, Rangel TFLVB, Bini LM. 2008. Model selection and information theory in
5 geographical ecology. *Global Ecology and Biogeography* 17: 479-488.
- 6 Dorado J, Vázquez DP. 2014. The diversity–stability relationship in floral production. *Oikos* 123:
7 1137–1143.
- 8 Ebeling A, Klein A-M, Schumacher J, Weisser WW, Tschardt T. 2008. How does plant richness
9 affect pollinator richness and temporal stability of flower visits? *Oikos* 117: 1808–1815.
- 10 Farr TG, Rosen PA, Caro E. et al. 2007. The shuttle radar topography mission. *Reviews of*
11 *Geophysics* 45: 1-33.
- 12 Figueiredo RA, Sazima M. 2000. Pollination Biology of Piperaceae species in Southeastern Brazil.
13 *Annals of Botany* 85: 455-460.
- 14 Fleming TH. 2005. The relationship between species richness of vertebrate mutualists and their food
15 plants in tropical and subtropical communities differs among hemispheres. *Oikos* 111: 556-
16 562.
- 17 Fonseca LCN, Vizentin-Bugoni J, Rech AR, 2015. Hummingbird-plant interactions and temporal
18 nectar availability in a restinga from Southeast Brazil. *Anais da Academia Brasileira de*
19 *Ciências* 87: 2163-2175.
- 20 Fontaine C, Dajoz I, Meriguet J, Loreau M. 2006. Functional Diversity of Plant–Pollinator Interaction
21 Webs Enhances the Persistence of Plant Communities. *PLoS Biol* 4: e1.
- 22 Friedman J, Barrett SCH. 2009. Wind of change, new insights on the ecology and evolution of
23 pollination and mating in wind-pollinated plants. *Annals of Botany* 103: 1515-1527.

- 1 Fründ J, Dormann CF, Holzschuh A, Tschardt T. 2013. Bee diversity effects on pollination depend
2 on functional complementarity and niche shifts. *Ecology* 94: 2042–2054.
- 3 Garcia RA, Cabeza M, Rahbek C, Araújo MB. 2014. Multiple dimensions of climate change and their
4 implications for biodiversity. *Science* 334: 6183.
- 5 Groom SVC, Stevens MI, Schwarz MP. 2014. Parallel responses of bees to Pleistocene climate
6 change in three isolated archipelagos of the southwestern Pacific. *Proceedings of the Royal
7 Society B* 281: 20133293.
- 8 Hansen MC, DeFries RS, Townshend JRG, Carroll M, Dimiceli C, Sohlberg R A. 2003. Global
9 Percent Tree Cover at a Spatial Resolution of 500 Meters: First Results of the MODIS
10 Vegetation Continuous Fields Algorithm. *Earth Interactions* 7: 1–15.
- 11 Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A. 2005. Very high resolution interpolated
12 climate surfaces for global land areas. *International Journal of Climatology* 25: 1965-1978.
- 13 Howe HF, Smallwood J 1982. Ecology of seed dispersal. *Annual Review of Ecology and Systematics*
14 13: 201–228.
- 15 Hu S, Dilcher DL, Taylor DW. 2012. Pollen evidence for the pollination biology of early flowering
16 plants. *Evolution of plant-pollinator relationships* (ed. by S. Patiny), pp. 165-236. Cambridge,
17 Cambridge University Press.
- 18 Jordano P. 2000. Fruits and frugivory. *Seeds: the ecology of regeneration in plant communities*(ed.
19 by M. Fenner), pp. 125–166. CAB International, Wallingford, UK.
- 20 Kearns CA, Inouye, DW, Waser NM. 1998. Endangered mutualisms: the conservation of plant–
21 pollinator interactions. *Annual Review Ecology Systematics* 29: 83–112.
- 22 Kissling WD, Eiserhardt WL, Baker WJ, Borchsenius F, Couvreur TLP, Balslev H, Svenning JC.
23 2012. Cenozoic imprints on the phylogenetic structure of palm species assemblages
24 worldwide. *Proceeding of the National Academy of Science* 109: 7379-7384.

- 1 Kissling WD, Rahbek C, Böhning-Gaese K. 2007. Food plant diversity as broad-scale determinant of
2 avian frugivore richness. *Proceedings of the Royal Society B* 274: 799-808.
- 3 Kühn I, Bierman SM, Durka W, Klotz S. 2006. Relating geographical variation in pollination types
4 to environmental and spatial factors using novel statistical methods. *New Phytologist* 172:
5 127-139.
- 6 Lever JJ, van Nes EH, Scheffer M, Bascompte J. 2014. The sudden collapse of pollinator
7 communities. *Ecology Letters* 17: 350-359.
- 8 Loarie SR, Dufy PB, Hamilton H, Asner GP, Field CB, Ackerly DD. 2009. The velocity of climate
9 change. *Nature* 462: 1052-1055.
- 10 Martín González AM, Dalsgaard B, Nogués-Bravo D, Graham CH, Schleuning M, Maruyama PK,
11 Abrahamczyk S, Alarcón R, Araujo AC, Araújo FP, Mendes de Azevedo S Jr, Baquero AC,
12 Cotton PA, Ingwersen TT, Kohler G, Lara C, Guedes Las-Casas FM, Machado AO, Machado
13 CG, Maglianesi MA, McGuire JA, Moura AC, Oliveira GM, Oliveira PE, Ornelas JF, Rodrigues
14 LdC, Rosero-Lasprilla L, Rui AM, Sazima M, Timmermann A, Varasin IG, Vizentin-Bugoni J,
15 Wang Z, Watts S, Rahbek C, Martinez ND. 2015. The macroecology of phylogenetically
16 structured hummingbird-plant networks. *Global Ecology and Biogeography* 24: 1212-1224.
- 17 McKinney AM, CaraDonna PJ, Inouye DW, Barr B, Bertelsen CD, Waser NM. 2012. Asynchronous
18 changes in phenology of migrating broad-tailed hummingbirds and their early-season nectar
19 resources. *Ecology* 93:1987-1993.
- 20 Niklas KJ. 1985. The aerodynamics of wind pollination. *Botanical Review* 51: 328-386.
- 21 Ollerton J, Johnson SD, Hingston AB. 2006. Geographical variation in diversity and specificity of
22 pollination systems. Pp. 283-308. *In*, Waser, N.M. and Ollerton, J. (eds.) *Plant-pollinator*
23 *interactions, from specialization to generalization*. University of Chicago Press, Chicago,
24 USA.

- 1 Ollerton J, Winfree R, Tarrant S. 2011. How many flowering plants are pollinated by animals? *Oikos*
2 120: 321-326.
- 3 Otto-Bliesner BL, Brady EC, Clauzet G, Tomas R, Levis S, Kothavala Z. 2006. Last Glacial
4 Maximum and Holocene Climate in CCSM3. *Journal of Climate* 19: 2526-2544.
- 5 R Development Core Team. 2014. R, A language and environment for statistical computing. R
6 Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org>.
- 7 Rangel TFLVB, Diniz-Filho JAF, Bini LM. 2010. SAM, a comprehensive application for Spatial
8 Analysis in Macroecology. *Ecography* 33: 46-50.
- 9 Rathcke B. 1983. Competition and facilitation among plants for pollination. *In*, Real, L. *Pollination*
10 *Biology*. Academic Press. pp. 305-325.
- 11 Regal PJ. 1982. Pollination by wind and animals: ecology of geographic patterns. *Annual Review of*
12 *Ecology and Systematics* 13: 497-524.
- 13 Sandel B, Arge L, Dalsgaard B, Davies RG, Gaston KJ, Sutherland WJ, Svenning JC. 2011. The
14 influence of late Quaternary climate-change velocity on species endemism. *Science* 334: 660-
15 664.
- 16 Santos T, Diniz-Filho JA, Rangel T, Bini LM. 2013. PVR: Computes phylogenetic eigenvectors
17 regression (PVR) and phylogenetic signal-representation curve (PSR) (with null and
18 Brownian expectations). R package version 0.2.1. <http://CRAN.R-project.org/package=PVR>
- 19 Schemske DW, Mittelbach GG, Cornell HV, Sobel JM, Roy K. 2009. Is there a latitudinal gradient
20 in the importance of biotic interactions? *Annual Review of Ecology, Evolution, and*
21 *Systematics* 40: 245-269.
- 22 Scherber C, Eisenhauer N, Weisser WW. et al. 2010. Bottom-up effects of plant diversity on
23 multitrophic interactions in a biodiversity experiment. *Nature* 468: 553-556.

- 1 Schleuning M, Ingmann L, Strauß R. et al. 2014. Ecological, historical and evolutionary
2 determinants of modularity in weighted seed-dispersal networks. *Ecology Letters* 17: 454-
3 463.
- 4 Svenning CJ, Skov F. 2007. Ice age legacies in the geographical distribution of tree species richness
5 in Europe. *Global Ecology and Biogeography* 16: 234-245.
- 6 Waser NM, Real LA. 1979. Effective mutualism between sequentially flowering plant species. *Nature*
7 281: 670-672.
- 8 Webb CO, Ackerly DD, Kembel SW. 2008. Phylocom: software for the analysis of phylogenetic
9 community structure and trait evolution. *Bioinformatics* 24: 2098-2100.
- 10 Webb CO, Donoghue MJ. 2005. Phylomatic: tree assembly for applied phylogenetics. *Molecular*
11 *Ecology Notes* 5: 181-183.
- 12 Whitehead DR. 1969. Wind-pollination in the angiosperms, evolutionary and environmental
13 considerations. *Evolution* 23: 28-35.
- 14 Willson MF, Irvine AK, Walsh NG. 1989. Vertebrate dispersal syndromes in some Australian and
15 New Zealand plant communities, with geographic comparisons. *Biotropica* 21: 133-147.
- 16 Wragg PD, Johnson SD. 2011. Transition from wind pollination to insect pollination in sedges,
17 experimental evidence and functional traits. *New Phytologist* 191: 1128-1140.
- 18 Yang CF, Wang QF, Guo YH. 2013. Pollination in a patchily distributed lousewort is facilitated by
19 presence of a co-flowering plant due to enhancement of quantity and quality of pollinator
20 visits. *Annals of Botany* 112: 1751-1758.

21

22 **Supporting information**

23 Appendix Table S1. Site descriptions

24 Appendix Table S2. Correlations between predictor variables

25 Appendix Table S3. Contemporary and historical determinants of the pollination mode

Table and Figures

Table 1. Correlations among the proportion of units in different taxonomic categories regarding pollination mode (animal *versus* wind pollination).

Taxa	Pearson correlation	t-value	Degrees of freedom	P-value
Species and Genera	0.971	31.2131	59	< 0.001
Species and Families	0.776	9.4789	59	< 0.001
Genera and Families	0.836	11.7435	59	< 0.001

Table 2. Contemporary and historical correlates (precipitation and temperature anomalies) of the proportion of animal-pollinated plant species in plant communities worldwide (n=67). The standardised regression coefficients are reported both for ordinary least square (OLS) and logistic regression, and reported for both an averaged model based on weighted w_i and minimum adequate models (MAMs), as in Diniz-Filho et al. (2008). For all MAMs based on OLS, we give AIC_c , Condition Number, Moran's I, and coefficients of determination (R^2). Finally, " $R^2_{\text{species richness}}$ ", " $R^2_{\text{topography}}$ ", " $R^2_{\text{current climate}}$ " and " $R^2_{\text{historical climate}}$ " reflect the unique variation explained by species richness, topography, current climate and historical climate, respectively. Note that historical climate stability is represented by temperature and precipitation anomaly between 21,000 years ago and the present (current minus Last Glacial Maximum - LGM precipitation/temperature, i.e. positive values reflect areas having been drier or colder at LGM than at present). See Appendix Table 2 for similar calculations when using temperature and precipitation velocities as historical climate stability measures.

	OLS			Logistic		
	Σw_i	Averaged	MAM [†]	Σw_i	Averaged	MAM [†]
Insularity	0.22	+0.02		0.96	-0.33	-0.33
Topography	0.84	+0.28	+0.28	1.00	+0.45	+0.42
Plant species richness	0.83	+0.23	+0.22	0.93	+0.24	+0.25

Open vegetation vs. forest	0.42	-0.22		1.00	+0.27	+0.29
MAP x Open vegetation vs forest	0.64	+0.37	+0.18	1.00	+1.18	+1.23
MAP (Mean Annual Precipitation)	0.25	-0.05		1.00	+0.13	+0.18
MAT (Mean Annual Temperature)	1.00	+0.69	+0.74	1.00	+1.42	+1.34
MAP seasonality	0.31	-0.11		0.48	-0.17	
MAT anomaly	0.70	+0.25	+0.25	1.00	+0.84	+0.85
MAP anomaly	0.24	+0.00		0.51	+0.18	
AIC (Akaike Information Criteria _c)			-73.29			668.9
Moran's Index			$\leq 0.08^{\text{NS}}$			
Condition Number			2.6			
R ²			0.49			
R ² _{species richness}			0.04			

$R^2_{\text{topography}}$	0.01
$R^2_{\text{current climate}}$	0.36
$R^2_{\text{historical climate}}$	0.02

** , $P < 0.01$; * , $P < 0.05$; ^{NS}, non-significant. † three models were equally fit (i.e. $\Delta\text{AIC}_c \leq 2$) containing the following variables, 1) Open vegetation vs forest, plant species richness, MAT, MAP x Open vegetation versus forest, topography, MAT anomaly; 2) plant species richness, MAT, topography, MAT anomaly; 3) Open vegetation vs forest, plant species richness, MAT, topography, MAT anomaly. † three other models were equally fit (i.e. $\Delta\text{AIC}_c \leq 2$) containing the following variables, 1) Open vegetation vs forest, insularity, plant species richness, MAT, MAP, MAP x Open vegetation versus forest, MAP seasonality, topography, MAT anomaly, MAP anomaly ; 2) Open vegetation vs forest, insularity, plant species richness, MAT, MAP, MAP x Open vegetation versus forest, MAP seasonality, topography, MAT anomaly; 3) Open vegetation vs forest, insularity, plant species richness, MAT, MAP, MAP x Open vegetation versus forest, topography, MAT anomaly, MAP anomaly.

Figure 1. Geographic patterns of animal pollination in 67 communities surveyed across the world. Fitted curves are second order polynomials; $r^2 = 0.28$, $F_{2, 47}$, 13.85, $P < 0.001$. On the map, dots were separated slightly within a small range to reduce the overlap of nearby sites.

Figure 2. Examples of species having traits representing wind (A-D) and animal (E-G) pollination used to classify plant species according to pollination mode in this study. A, *Ambrosia artemisiifolia* L. (Asteraceae) a diclinous wind pollinated plant with minute flowers that produce a large amount of pollen (Photo by Andrew Butko via Wiki images), B, *Mercurialis annua* L. (Euphorbiaceae) a dioecious wind-pollinated plant species; the circle highlights the female flower and, in detail at the right side of the image, we show the inflorescence of a male individual (Photo by Hasan Yldirin), C, *Olyra ciliatifolia* Raddi (Poaceae) female flower with feathery stigmas (Photo by Pedro Viana), D, *Paspalum notatum* Flügge (Poaceae) showing the flexible white filaments; the arrow is pointing to the wide anther aperture (Photo by Pedro Viana), E, *Turnera ulmifolia* L. (Passifloraceae) a 3D heterostyle species and a butterfly, one of its animal pollinators, F, *Couroupita guianensis* Aubl. (Lecythidaceae) with food (colourful) and pollination (white) specialised stamens (Photo by Hipolito Neto), G, *Gongora bufonia* Lindl. (Orchidaceae) being pollinated by a male euglossine bee (*Eufrisea violacea*) when collecting scented oil from its labellum (Photo by Carlos Eduardo Coquinho).

Figure 3. The relationship between the proportion of animal pollinated plant species per community and the most important and consistent predictors, a) plant richness; b) temperature; c) precipitation. Open symbols illustrate open vegetation types whereas filled symbols

illustrate forest communities. All relationships significant in simple univariate correlations are shown: all vegetation types ($n = 67$; dashed line); open vegetation types ($n = 51$, dotted line); and forest ($n = 16$, full line). See Table 2 and Tables S1-S2 for statistics.

Appendix Table S1. Site descriptions for each community analysed in the study: “The macroecology of animal *versus* wind pollination”. Each number in the first column corresponds to a reference below.

N°	Study site	Mainland = 0	Forest: 1	Total	N° wind	% animal	Latitude	Longitude
		Island = 1	Other = 0	plants:	pol. species	pollination	Decimalised	Decimalised
1	Juan Fernandez Islands, Chile	1	0	149	70	53.02	-33.64	-78.84
2	Coastal scrub, UK	1	0	37	9	75.68	54.40	-0.51
3	Degraded tropical scrub/secondary	0	1	82	3	96.34	22.28	114.2
4	Serra da Bocaina - Grassland	0	0	179	55	69.27	-22.73	-44.61
5	Cerrado, Brazil	0	0	294	52	82.31	-22.87	-48.49
6	semi-arid scrub - Spain	0	0	31	6	80.65	37.01	-6.55
7	Amami Island, Japan	1	0	103	6	94.17	28.32	129.42
8	All habitats across New Caledonia	1	0	97	3	96.91	-21.17	165.25
9	La Selva, Costa Rica	0	1	283	13	95.41	10.45	-84.00
10	Caatinga, Brazil	0	0	147	3	97.96	-8.45	-36.76
11	various localities, Faroe Islands	1	0	70	19	72.86	62.00	6.783
12	Guyana Highlands, Venezuela	0	0	55	6	89.09	5.58	-61.71
13	Rainforest, Malaysia	1	1	262	0	100.00	4.20	114.5
14	Alpine pavement plain, California	0	0	17	4	76.47	34.30	-116.86
15	Woodland, USA	0	1	208	57	72.60	42.1	-111.59
16	various localities, St Kilda	1	0	92	39	57.61	57.81	-8.58
17	Palm swamp, Venezuela	0	1	33	5	84.85	8.93	-67.25
18	Prairie, Illinois	0	0	409	14	96.58	39.46	-89.90
19	Flooded/Upland Rainforest, Colombia	0	1	90	2	98.51	-0.61	-72.33
20	Lower alpine, Scotland	1	0	153	8	94.77	56.50	-3.11
21	Serra do Cipo	0	0	64	5	92.19	-23.336667	-45.14
22	Botucatu	0	0	34	3	91.18	-22.844167	-48.68
23	Santa Virginia	0	1	70	8	88.57	-19.334722	-43.58

24	Picinguaba - restinga	0	0	47	5	89.36	-23.358056	-44.85
25	Picinguaba - Atlantic forest	0	1	43	2	95.35	-23.338611	-44.83
26	La Floresta - Canelones - Uruguay -	0	0	24	10	58.33	-34.760706	-55.69
27	Quebrada de los Cuervos - Uruguay -	0	0	42	14	66.67	-32.935091	-54.46
28	Kumu - Guyana - rainforest	0	1	59	3	94.92	3.266667	-59.75
29	Kumu - Guyana - savannah	0	0	43	6	86.05	3.266667	-59.77
30	Wahroonga - South Africa	0	0	73	4	94.52	-29.616667	30.13
31	Mantanay - Peru	0	0	148	6	95.95	-13.2	-72.08
32	Scrub Field - Northampton - UK	1	0	162	35	78.40	52.26	-0.88
33	Bahia de Patano - Venezuela	0	0	78	12	84.62	10.466667	-67.75
34	Guimar Badlands, Tenerife	1	0	18	2	88.89	28.33	-16.42
35	Mean JasperRidge Plant Covariates -	0	0	109	24	78.98	37.417	-122.19
36	Mean JasperRidge Plant Covariates -	0	1	78	16	77.85	37.417	-122.19
37	JasperRidge Plant Covariates -	1	0	16	7	56.25	37.701067	-123.00
38	MatherPlantCovariatesPUPPLEEsEt -	0	0	92	31	66.30	37.8774	-119.25
39	MatherPlantCovariatesPUPPLEEsEt -	0	0	58	6	89.66	37.8374	-120.30
40	MatherPlantCovariatesPUPPLEEsEt -	0	1	132	19	85.61	37.8774	-119.29
41	MatherPlantCovariatesPUPPLEEsEt -	0	0	134	33	75.37	38.0095	-123.00
42	TimberlinePlantDataPurple - Dore	0	0	60	19	68.33	37.9683	-119.30
43	TimberlinePlantDataPurple -	0	0	95	47	50.53	37.9382	-119.25
44	TimberlinePlantDataPurple -	0	1	91	34	62.64	37.9612	-119.29
45	TimberlinePlantDataPurple - Talus	0	0	126	38	69.84	37.9462	-119.25
46	Virginia Basin (Colorado)	0	0	64	6	90.63	38.98	-106.966667
47	Mean Hangklip, South Africa	0	0	124	41	65.55	-34.25	18.75
48	Brattnesdalen, Norway	0	0	18	8	55.56	70.25	22.07
49	Fjorddalen, Norway	0	0	31	10	67.74	70.19	22.10
50	various localities, Australia	0	1	148	8	94.59	-19.18	146.75
51	Torrey Pines - California	0	0	85	21	75.29	32.89	-117.24
52	Japatul Valley - California	0	0	91	15	83.52	32.78	-116.68
53	Echo Valley - California	0	0	56	4	92.86	32.89	-116.65

54	Mount Laguna - California	0	1	65	15	76.92	32.86	-116.41
55	Ocotillo - California	0	0	80	14	82.50	32.74	-115.99
56	Papudo - Chile	0	0	95	23	75.79	-32.55	-71.44
57	Fundo Santa Laura - Chile	0	0	106	18	83.02	-33.26	-70.85
58	Cerro Potrerillo - Chile	0	0	47	11	76.60	-30.29	-70.57
59	El Tofo - Chile	0	0	44	5	88.64	-29.88	-71.20
60	Alpine - montane Colorado	0	0	68	17	75.00	38.68	-107.115532
61	Aspen - montane Colorado	0	1	56	15	73.21	38.73	-106.772653
62	Sage - montane Colorado	0	0	45	9	80.00	38.73	-106.823376
63	Grassland - montane Colorado	0	0	103	24	76.70	38.95	-106.988824
64	Spruce-fir - montane Colorado	0	1	50	11	78.00	38.86	-107.101081
65	Salt marsh, Canada	0	0	18	6	66.67	49.08	-125.85
66	Sphagnum bog, Canada	0	0	33	13	60.61	49.08	-125.86
67	Subalpine meadow, Canada	0	0	45	11	75.56	49.11	-120.84

Appendix Table S1 - References

- 1- Bernadello, G. et al. 2001. A survey of floral traits, breeding systems, floral visitors, and pollination systems of the angiosperms of the Juan Fernandez Islands (Chile). - *Bot. Rev.* 67: 255 - 308.
- 2- Burkill, I. H. 1897. Fertilization of some spring flowers on the Yorkshire coast. - *J. Bot.* 35: 92 - 189.
- 3- Corlett, R. T. 2001. Pollination in a degraded tropical landscape: a Hong Kong case study. - *J. Trop. Ecol.* 17: 155-161.
- 4- Freitas, L. and Sazima, M. 2006. Pollination biology in a tropical high-altitude grassland in Brazil: interactions at the community level. *Ann.Missouri Bot.Gard.* 93: 465-516.
- 4- Martinelli, G. 1989. *Campos de Altitude*. Editora Index, Rio de Janeiro.
- 5- Gottsberger, G. and Silberbauer-Gottsberger, I. 2006. *Life in the Cerrado*. Reta Verlag.
- 6- Herrera, J. 1988. Pollination relationships in southern Spanish mediterranean shrublands. *J. Ecol.* 76: 274 - 287.
- 7- Kato, M. 2000. Anthophilous insect community and plant-pollinator interactions on Amami Islands in the Ryukyu Archipelago, Japan. - *Contrib. Biol. Lab. Kyoto Univ.* 29: 157 – 252.
- 8- Kato, M. and Kawakita, A. 2004. Plant-pollinator interactions in New Caledonia influenced by introduced honey bees. *Am. J. Bot.* 91: 1814-1827.

- 9- Kress, W. J. and Beach, J. H. 1994. Flowering plant reproductive systems. - *In*: McDade, L. A. et al. (eds), *La Selva: ecology and natural history of a Neotropical rainforest*. Univ. Chicago Press, pp. 161 - 182.
- 10- Machado, I. C. and Lopes, A. V. 2004. Floral traits and pollination systems in the caatinga, a Brazilian tropical dry forest. *Ann. Bot.* 94: 365-376.
- 11 - 12- Ollerton, J. Unpublished data, available upon request from the author via Jeff.Ollerton@northampton.ac.uk
- 13- Momose, K. et al. 1998. Pollination biology in a lowland dipterocarp forest in Sarawak, Malaysia. Characteristics of the plant-pollinator community in a lowland dipterocarp forest. *Am. J. Bot.* 85: 1477-1501.
- 14- O'Brien, M. H. 1980. The pollination biology of a pavement plain: pollinator visitation patterns. - *Oecologia* 47: 213-218
- 15- Ostler, W. K. and Harper, K. T. 1978. Floral ecology in relation to plant species diversity in the Wasatch Mountains of Utah and Idaho. *Ecology* 59:848-861.
- 16 and 48-49- Holmes, N. 2014. Unpublished data, available upon request from the author via Naomi.Holmes@northampton.ac.uk
- 17- Ramirez, N. and Brito, Y. 1992. Pollination biology in a palm swamp community in the Venezuelan Central Plains. *Bot. J. Linn. Soc.* 110: 277-302.

- 18- Robertson, C. 1928. Flowers and insects: lists of visitors of four hundred and fifty-three flowers. Privately published, Carlinville, Illinois.
- 19- van Dulmen, A. 2001. Pollination and phenology of flowers in the canopy of two contrasting rain forest types in Amazonia, Colombia. - Plant Ecol. 153: 73-85.
- 20- Willis, J. C. and Burkill, I. H. 1895. Flowers and insects in Great Britain. Part I. Ann. Bot. 9: 227-273.
- 20- Willis, J. C. and Burkill, I. H. 1903a. Flowers and insects in Great Britain. Part II. Ann. Bot. 17: 313-349
- 20- Willis, J. C. and Burkill, I. H. 1903b. Flowers and insects in Great Britain. Part III. Ann. Bot. 17: 539-570.
- 20- Willis, J. C. and Burkill, I. H. 1908. Flowers and insects in Great Britain. Part IV. Ann. Bot. 22: 603-649.
- 21-25- Ollerton, J. and Rech, A.R. 2013. unpublished data, available upon request from the authors via Jeff.Ollerton@northampton.ac.uk
- 26-27- Rech, A.R. 2014. unpublished data, available upon request from the author via andrerodrigorech@gmail.com
- 28-34- Ollerton, J. Unpublished data, available upon request from the author via Jeff.Ollerton@northampton.ac.uk
- 35-36- Moldenke, D. Unpublished data, requested from the author.
- 37-46- Moldenke, D. Unpublished data, requested from the author.

- 47 and 50- Ollerton, J. Unpublished data, available upon request from the author via Jeff.Ollerton@northampton.ac.uk
- 51-59- Moldenke, A. R. 1979. Pollination ecology as an assay for ecosystemic organization: convergent evolution in Chile and California. *Phytologia* 42: 415-454.
- 60-64- Moldenke, A. R. and Lincoln, P. G. 1979. Pollination ecology in montane Colorado: a community analysis. *Phytologia* 42: 349 - 379.
- 65-67- Pojar, J. 1974. Reproductive dynamics of four plant communities of southwestern British Columbia. *Can. J. Bot.* 52: 1819 - 1834.

Appendix Table S2. Correlations between the proportion of animal pollinated plant species and all predictor variables for all vegetation types (n = 67, except for velocities which had one data point less as we were unable to extract it for St Kilda Island, Scotland) above the diagonal, and separately for open vegetation types (n = 51, except for velocities which had one data point less as we were unable to extract it for St Kilda Island, Scotland) below the diagonal.

	% animal pollinated	Plant richness	MAT	MAP	MAP seasonality	Topography	MAT anomaly	MAP anomaly	MAT velocity	MAP velocity
% animal pollinated		+0.32**	+0.57*	+0.29[†]	+0.09 ^{NS}	-0.11 ^{NS}	-0.14 ^{NS}	+0.03 ^{NS}	+0.07 ^{NS}	+0.13 ^{NS}
Plant richness	+0.28*		+0.11 ^{NS}	+0.11 ^{NS}	-0.05 ^{NS}	-0.11 ^{NS}	+0.06 ^{NS}	+0.23 ^{NS}	+0.16 ^{NS}	+0.05 ^{NS}
MAT (Mean Annual Temperature)	+0.42*	+0.09 ^{NS}		+0.44*	+0.35*	-0.45**	-0.50*	+0.01 ^{NS}	+0.05 ^{NS}	+0.37**
MAP (Mean Annual Precipitation)	-0.02 ^{NS}	-0.04 ^{NS}	+0.24 ^{NS}		-0.17 ^{NS}	-0.25[†]	-0.20 ^{NS}	+0.33**	+0.14 ^{NS}	+0.25*
MAP seasonality	+0.14 ^{NS}	+0.02 ^{NS}	+0.39*	-0.18*		+0.12 ^{NS}	-0.53*	-0.05 ^{NS}	-0.36[†]	+0.20 ^{NS}
Topography	+0.04 ^{NS}	-0.08 ^{NS}	-0.36**	-0.17 ^{NS}	+0.19 ^{NS}		-0.04 ^{NS}	-0.16 ^{NS}	-0.71**	-0.47**
MAT anomaly	-0.08 ^{NS}	+0.12 ^{NS}	-0.51*	-0.10 ^{NS}	-0.60**	-0.13 ^{NS}		-0.17 ^{NS}	+0.50**	-0.21 ^{NS}
MAP anomaly	-0.22 ^{NS}	+0.08 ^{NS}	-0.17 ^{NS}	+0.22 ^{NS}	+0.06 ^{NS}	-0.01 ^{NS}	-0.13 ^{NS}		+0.16 ^{NS}	+0.10 ^{NS}
MAT velocity	+0.01 ^{NS}	+0.16 ^{NS}	-0.04 ^{NS}	+0.10 ^{NS}	-0.41*	-0.70**	+0.58**	+0.05 ^{NS}		+0.40*
MAP velocity	+0.02 ^{NS}	-0.01 ^{NS}	+0.29**	+0.14 ^{NS}	+0.17 ^{NS}	-0.42**	-0.15 ^{NS}	-0.05 ^{NS}	+0.43*	

** $P < 0.01$; * $P < 0.05$ when P-values based on degrees of freedom corrected for spatial autocorrelation using Dutilleul's (1993) method;

†significant when using traditional non-spatial statistics, but non-significant when corrected for spatial autocorrelation; ^{NS}non-significant.

Appendix Table S2, continued. Correlations between predictor variables separately for forest (n = 16).

	% animal pollinated	Plant richness	MAT	MAP	MAP seasonality	Topography	MAT anomaly	MAP anomaly	MAT velocity	MAP velocity
% animal pol.		+0.30 ^{NS}	+0.82[†]	+0.84[*]	-0.01 ^{NS}	-0.59[†]	-0.46 ^{NS}	+0.37 ^{NS}	+0.32 ^{NS}	+0.50[†]
Plant richness			+0.05 ^{NS}	+0.38 ^{NS}	-0.29 ^{NS}	-0.22 ^{NS}	-0.29 ^{NS}	+0.57[*]	+0.21 ^{NS}	+0.26 ^{NS}
MAT (Mean Annual Temperature)				+0.74 [†]	+0.32 ^{NS}	-0.73[†]	-0.67[†]	+0.19 ^{NS}	+0.34 ^{NS}	+0.58[†]
MAP (Mean Annual Precipitation)					-0.13 ^{NS}	-0.52[†]	-0.70[*]	+0.42 ^{NS}	+0.33 ^{NS}	+0.58[†]
MAP seasonality						-0.14 ^{NS}	-0.26 ^{NS}	-0.30 ^{NS}	-0.13 ^{NS}	+0.26 ^{NS}
Topography							+0.55 [†]	-0.56[*]	-0.78[*]	-0.61[†]
MAT anomaly								-0.41 ^{NS}	-0.17 ^{NS}	-0.64[*]
MAP anomaly									+0.56[*]	+0.44 ^{NS}
MAT velocity										+0.36 ^{NS}
MAP velocity										

**P < 0.01; *P < 0.05 when P-values based on degrees of freedom corrected for spatial autocorrelation using Dutilleul's (1993) method;

†significant when using traditional non-spatial statistics, but non-significant when corrected for spatial autocorrelation; ^{NS}non-significant.

Appendix Table S3. Contemporary and historical determinants (precipitation and temperature velocities) of the proportion of animal-pollinated plant species in plant communities worldwide (n=66). The standardized regression coefficients are reported both for ordinary least square (OLS)

and logistic regression, and reported for both an averaged model based on weighted w_i and minimum adequate models (MAMs), as in Diniz-Filho *et al.* (2008). For all MAMs based on OLS, we give AIC_c, Condition Number, Moran's I, and coefficients of determination (R^2). Finally, " $R^2_{\text{species richness}}$ ", " $R^2_{\text{current climate}}$ " and " $R^2_{\text{historical climate}}$ " reflect the unique variation explained by species richness, current climate and historical climate, respectively. Notice that historical climate stability is represented by temperature and precipitation velocity between 21000 years ago and now, and that topography is not included as strongly correlated with velocities. See Table 2 for similar calculations using precipitation and temperature anomalies as historical climate stability measures.

	OLS			Logistic		
	Σw_i	Averaged	MAM [†]	Σw_i	Averaged	MAM
Insularity	0.25	+0.05		0.75	-0.24	-0.25
Plant species richness	0.85	+0.24	+0.24	0.85	+0.21	+0.20
Open vegetation vs forest	0.44	-0.24		1.00	+0.31	+0.29
MAP x Open vegetation vs forest	0.67	+0.39	+0.20	1.00	+1.22	+1.23
MAP (Mean Annual Precipitation)	0.27	-0.07		1.00	+0.03	+0.02
MAT (Mean Annual Temperature)	1.00	+0.50	+0.47	1.00	+0.82	+0.84

MAP seasonality	0.31	-0.10		0.87	-0.26	-0.27
MAT velocity	0.23	-0.00		1.00	+0.42	+0.40
MAP velocity	0.29	-0.09		0.22	-0.01	
Akaike Information Criteria _c			-69.94			675.9
Moran's Index			≤0.14*			
Condition Number			1.5			
R ²			0.43			
R ² _{species richness}			0.05			
R ² _{current climate}			0.32			
R ² _{historical climate}			0.00			

**P < 0.01; *P < 0.05; ^{NS}non-significant. † five other models were equally fit (i.e. $\Delta AIC_c \leq 2$) containing the following variables, 1) Open vegetation vs forest, plant species richness, MAT, MAP x Open vegetation versus forest; 2) plant species richness, MAT; 3) plant species

richness, MAT, MAP x Open vegetation versus forest, MAP velocity; 4) plant species richness, MAT, MAP x Open vegetation versus forest, MAP seasonality; 5) Open vegetation vs forest, plant species richness, MAT.